

# UC Irvine

## UC Irvine Previously Published Works

### Title

Bunyaviruses are Dependent on K2p Channels to Infect Cells

### Permalink

<https://escholarship.org/uc/item/43b5m7k0>

### Journal

Biophysical Journal, 114(3)

### ISSN

0006-3495

### Authors

Hover, Samantha  
Barr, John N  
AN Goldstein, Steve  
et al.

### Publication Date

2018-02-01

### DOI

10.1016/j.bpj.2017.11.2684

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

**2418-Pos Board B434**

**Bunyaviruses are Dependent on K<sub>2p</sub> Channels to Infect Cells**

**Samantha Hover<sup>1</sup>**, John N. Barr<sup>1</sup>, Steve AN Goldstein<sup>2</sup>, Jamel Mankouri<sup>1</sup>.

University of Leeds, Leeds, United Kingdom, <sup>2</sup>Loyola University Chicago, Chicago, IL, USA.

All viruses must infect cells in order to multiply and cause disease. Once inside cells, viruses hijack normal cellular processes to create an environment that favours their own survival, typically at the expense of the host. Through understanding how viruses do this, we can design new strategies to prevent viruses from multiplying, and thus stop their ability to cause disease. The *Bunyaviridae* are the largest family of negative stranded RNA viruses. These are a group of over 350 different viruses spread throughout the globe. They are predominantly transmitted to humans by insects and are capable of causing fatal disease in humans, often as a result of devastating hemorrhagic fevers. Bunyaviruses are a serious and worrying threat to human health because they have enormous capacity to mutate and evolve into new strains. In addition, the fact that bunyaviruses are mostly spread by biting insects that are highly mobile means they can rapidly move into new geographic locations, causing widespread disease. Despite this huge threat, no preventative or therapeutic measures currently exist for any bunyavirus-mediated disease. Using model viruses within the *Bunyaviridae* family, predominantly Bunyamwera virus (BUNV), we previously identified cellular potassium (K<sup>+</sup>) channels as an essential host cell factor required for bunyavirus infection. Using a rationale panel of K<sub>p</sub> channel modulators, we identified the two-pore K<sub>p</sub> (K2P) channel family as those required by BUNV. We have identified four possible K2P channels we believe are facilitating a cellular process required during BUNV infection. Using genetic silencing we plan to determine the specific channel(s) involved and from this, identify a new drug target for the development of novel anti-bunyavirus therapies.